

<u>Cabinets (LFBSC), Major Spills – Class 2 and 3 Organisms.....</u>	<u>27</u>
<u>Liquid Disinfectants.....</u>	<u>28</u>
<u>VSU Biosafety Forms.....</u>	<u>30</u>
· <u>Memorandum of Understanding and Agreement (MUA) for Recombinant DNA Experiments.....</u>	<u>31</u>
· <u>MUA for Biohazards other than Recombinant DNA Experiments.....</u>	<u>34</u>
· <u>VSU Biosafety Infectious Agent Risk Assessment.....</u>	<u>37</u>
· <u>Laboratory Equipment Approval Form.....</u>	<u>42</u>
· <u>Biohazardous Research Checklist.....</u>	<u>43</u>

I. Introduction

In conjunction with the U.S. Patriot Act and Valdosta State University’s Campus Homeland Security (www.valdosta.edu/legal/chs), the following policy outlines the practices embodied in the aforementioned authority as they relate to selected agents/chemicals. This policy defines the responsibilities of certain individuals, as well as protocol/practices. Its purpose is to provide guidelines for safe handling of biohazardous materials (including biological agents, toxins, and recombinant DNA) by employees and students at Valdosta State University. These guidelines are offered for the protection of students and employees at Valdosta State University as well as the general community.

II. Primary Policy and Procedures Source for VSU Biosafety Manual Development

This Biosafety Manual is based largely on the Biosafety Manual of the University of Georgia and the relative contents contained therein are used with expressed permission. Biosafety policies and procedures at Valdosta State University conform to policies and procedures as prescribed by the Centers for Disease Control and the National Institute of Health and published in Biosafety in Microbiological and Biomedical Laboratories, CDC/NIH, Fourth Edition, U.S. Department of Health and Human Services, U.S. Government Printing Office, May 1999. (Document is available at

<http://www.cdc.gov/od/ohs/biosfty/bmbl4/bmbl4toc.htm>).

III. Resources on Biosafety

Many excellent resources are available on various aspects of biosafety; several are listed below.

- Office of Health and Safety Information System, CDC.
(<http://www.cdc.gov/od/ohs/default.htm>)
- CDC Office of Health and Safety, Biosafety Documents
<http://www.cdc.gov/od/ohs/biosfty/biosfty.htm>
- Biosafety Resources on the Internet, American Biological Safety Association
(<http://www.absa.org/resources/resource.htm>).
- American Society for Microbiology, Resources Related To Biological Weapons Control And Bioterrorism Preparedness <http://www.asmta.org/pasrc/bioprep.htm>
- *Biosafety in Microbiological and Biomedical Laboratories, CDC/NIH*, Fourth Edition, U.S. Department of Health and Human Services, U.S. Government Printing Office, May 1999. (available at <http://www.cdc.gov/od/ohs/biosfty/bmbl4/bmbl4toc.htm>).
- Risk Group Classification for Infectious Agents (available at the web site of the American Biological Safety Association, <http://www.absa.org/riskgroups/default.htm>).
- Bloodborne pathogens (29CFR, 1910.1030), Occupational Safety and Health Administration (available at http://www.osha-slc.gov/OshStd_data/1910_1030.html)
- *Primary Containment for Biohazards: Selection, Installation and Use of Biological Safety Cabinets*, U.S. Dept. of Health and Human Services, September 1995. (available at <http://www.cdc.gov/od/ohs/biosfty/bsc/bsc.htm>).
- Interstate Shipment of Etiologic Agents (42CFR Part 72), Federal Register, July 21, 1980. (available at <http://www.cdc.gov/od/ohs/biosfty/shipregs.htm>).
- Additional requirements for facilities transferring or receiving select agents; Final Rule (42 CFR Part 72.6), Federal Register, Oct. 24, 1996. (available at <http://www.cdc.gov/od/ohs/lrsat/regmat.htm>).
- Select Agent Rule <http://www.phppo.cdc.gov/nltn/sar.asp>

- Guidelines for the Safe Transport of Infectious Substances and Diagnostic Specimens, World Health Organization, 1997. (available at <http://www.absa.org/resources/who-97.pdf>).
- Notification of Possession of Select Agents or High Consequence Livestock Pathogens and Toxins. Federal Register Vol 67, No. 151, Tuesday, August 6, 2002
- Agricultural Bioterrorism Protection Act: Biological agents and toxins; possession. Federal Register Vol. 67, No. 155, Monday, August 12, 2002
- USDA Animal and Plant Health Inspection Service, Veterinary Services, National Center for Import and Export <http://www.aphis.usda.gov/vs/ncie/>
- USDA Animal and Plant Health Inspection Service, Plant Protection and Quarantine <http://www.aphis.usda.gov/ppq/>
- Agricultural Permits, U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine, (<http://www.aphis.usda.gov/ppq/permits/index.html>)
- U.S. Regulatory Oversight in Biotechnology, U.S. Department of Agriculture, U.S. Environmental Protection Agency, and U.S. Food and Drug Administration (Unified Homepage) (<http://www.aphis.usda.gov/biotech/OECD/usregs.htm#fdalaw>)
- *Guidelines for Research Involving Recombinant DNA Molecules (NIH Guidelines)*, April 2002. (available at http://www4.od.nih.gov/oba/rac/guidelines_02/NIH_Guidelines_Apr_02.htm)
- Code of Federal Regulations (U.S. Government) <http://www.access.gpo.gov/nara/cfr/index.html>
- Federal Register (1995-present) http://www.gpo.gov/su_docs/aces/aces140.html
- U.S. Congress, 1988. *Two Edera*

on protecting workers in the immediate area; whereas, secondary containment deals with protection of the environment and people outside the immediate area. Containment is accomplished through: (1) the use of appropriate procedures, (2) the use of safety equipment, and (3) conducting work in an appropriately designed facility.

V. Responsibilities

A. Faculty and Professional Staff (Principal

1. Report in writing to the Department Head and Biosafety Officer any laboratory or field accident, exposure of personnel, suspected illness, escape from containment of biohazardous agents, and significant problems pertaining to the operation and

on biohazards for the University community.

4. Review periodically biohazardous research being conducted at the University to insure that the requirements of the University, funding sources, and regulatory agencies are being fulfilled.
5. Recommend to the University Administration appropriate sanctions for non-compliance with biosafety standards, guidelines, or regulations.
- 6.

1. developing and maintaining appropriate policies regarding the conduct of potentially biohazardous research, education, and service activities.
2. developing mechanisms for insuring adherence to biosafety policies.
3. providing the resources necessary for the construction of safe research and teaching facilities and for the implementation of the biosafety program.
4. providing adequate resources for the dissemination of information on biohazards and biosafety procedures, including training programs and workshops.
5. providing resources for appropriate medical surveillance measures to protect the health and safety of employees.
6. providing appropriate and sufficient legal protection for faculty and staff members who conduct activities in compliance with appropriate regulations and guidelines.

VI. Biohazardous Research

-

A. General Definitions

1. Institutional Biosafety Committee (IBC) - The University Committee appointed by the President of Valdosta State University and which meets the requirements specified by NIH in its "Guidelines for Research Involving Recombinant DNA Molecules". The Committee also reviews, approves, activities involving recombinant DNA as well as other biohazards identified in this manual.
2. Biohazard - infectious agents, or parts or products thereof, presenting a real or potential risk to the well-being of humans, other animals, or plants directly through infection and/or toxicity or indirectly through disruption of the environment; and venomous vertebrate or invertebrate animals and other toxic organisms presenting a real or potential risk to humans.
3. Class 2 Agents - are those agents that are to be handled using Biosafety Level 2 or greater containment facilities and practices.
4. Class 3 Agents - are those agents that are to be handled using Biosafety Level 3 or greater containment facilities and practices.
5. Class 4 Agents - are those agents that are to be handled using Biosafety Level 4 containment facilities and practices.
6. Genetic Engineering - the genetic modification of organisms by recombinant DNA techniques.
7. Plant Pests - Any living stage (including active and dormant forms) of insects, mites, nematodes, slugs, snails, protozoa, or other invertebrate animals, bacteria, fungi, other parasitic plants or reproductive parts thereof; viruses; or any organisms similar to or allied with any of the foregoing; or any infectious agents or substances, which can directly or indirectly injure or cause disease or damage in or to any plants or parts thereof, or any processed, manufactured, or other products of plants. (USDA - 7 CFR 340.1)
8. Regulated Article - any organism which has been altered or produced through genetic engineering, if the donor organism, recipient organism, or vector or vector agent belongs to any genera or taxa designated in 7 CFR 340.2 and meets the definition of plant pest, or

Pathogens", Federal Register, 56:235:64004-64174, December 6, 1991.

· *Biosafety in Microbiological and Biomedical Laboratories, CDC/NIH*, Fourth Edition, U.S. Department of Health and Human Services, U.S. Government Printing Office, May 1999. (available at

Reference:

their article "Microbiological Safety Evaluation of an Industrial Refuse Incinerator" (Applied Microbiology 16:2:291-95) reported on various times required for autoclaving selected animal carcasses, animal bedding materials, and eggs. W

BSL 1 Standard Microbiological Safety Practices

1. Access to the laboratory is limited or restricted at the discretion of the laboratory director. Limiting access to control the in/out traffic when experiments are in progress reduces sources of distraction and disturbance which may result in accidents. Closing laboratory doors during experiments is one method of controlling in/out traffic. This also allows for the exclusion of special category persons (children, immunosuppressed persons, etc.) during times of potential exposure.
2. Work surfaces are decontaminated at least once a day and following any spill of viable material. Contaminated equipment must be decontaminated according to any local, state or federal regulations before it is sent for repair or maintenance or packaged for transport or surplused. This practice assists in the control of general contamination of the laboratory and reduces infection potential among laboratory personnel as well as repair people and others in contact with with laboratory equipment.
3. All contaminated liquid and solid wastes are decontaminated prior to disposal. Disposal of biomedical wastes shall be accomplished so as to comply with state and federal laws and regulations, see Part VI.
4. Mechanical pipetting devices are used; mouth pipetting is prohibited. Mechanical pipetting is easy and accurate, and prevents the ingestion of the materials being pipetted. Several older publications referred to human infection and death associated with the mouth pipetting of pathogens.
5. Eating, drinking, smoking, and the application of cosmetics are not permitted in the work area. Food may be stored cabinets and refrigerators designated and used for this purpose only. Food storage cabinets and refrigerators should be located outside the work area. Storage and/or consumption of food and drink and application of cosmetics in biohazardous work areas may result in exposure to laboratory personnel via the contamination of these products.
6. Personnel wash their hands after they handle viable materials and animals, after removing gloves, and before leaving the laboratory or animal facility. This practice reduces the potential for ingestion and/or absorption of harmful microorganisms and hazardous chemicals. It also reduces the likelihood of exposure to hazardous chemicals.

at BSL 1.

Biosafety Level 2 / Laboratory

Biosafety Level 2 is suitable for work involving agents of moderate potential hazard to personnel and the environment (including plants and other animals). The practices, equipment, and laboratory design are appropriate for clinical, diagnostic, teaching, and basic research with a broad spectrum of indigenous moderate-risk agents associated with human disease and/or which may negatively impact the environment. Laboratory procedures which generate aerosols may increase the risk and therefore are to be conducted in a biological safety cabinet and/or other primary containment equipment.

Biosafety Level 2 facilities and procedures are those that are basic in a good quality laboratory working with microorganisms, genetic materials, cell/tissue cultures, and carcinogens. In addition to the BSL 1 Standard Microbiological Safety Procedures, the following Special Practices are implemented:

1. Access to the laboratory is limited or restricted by the laboratory supervisor when work with biohazardous agents is in progress. The laboratory director has the final responsibility for assessing each circumstance and determining who may enter or work in the laboratory or animal rooms. Keeping laboratory doors closed during experiments is recommended. Persons who are at increased risk of acquiring infection or for whom infection may be unusually hazardous are not allowed in the laboratory or animal rooms. For example, persons who are immunocompromised or immunosuppressed may be at risk of acquiring infections.
2. The principal investigator is responsible for providing training of laboratory personnel in the potential hazards and safety procedures. Knowledgeable personnel work more efficiently and effectively in the laboratory by reducing the risks of accidents that could result in personal injury or loss of research effort. Georgia Law "Public Employee Hazardous Chemical Protection and Right to Know Act of 1988" (The Official Code of Georgia Annotated Title 45 Chapter 22) and the Department of Labor regulations (Chapter 300-3-19) provide for training of employees using hazardous chemicals. It only makes sense that investigatory also provide training to employees using biohazards.
3. When research involves working with or storing biohazardous agents in the laboratory a hazard warning sign incorporating the universal biohazard symbol is posted on the access door. The principal investigator is ultimately responsible for informing persons, including emergency personnel, of any special requirement for entering the laboratory.
4. Before leaving the laboratory areas, protective clothing (lab coats, aprons, etc.) is removed and left in the laboratory. This practice helps prevent infectious agents from being carried from the laboratory on contaminated clothing.
5. Animals not involved in the work being performed are not permitted in the laboratory. Pets or other noninvolved animals may bring unwanted organisms into the laboratory or may carry infectious agents from the laboratory into the home, into other areas of the building, or into the community.
6. Special care is taken to avoid contamination of skin and mucous membranes with infectious materials; appropriate personal protective equipment (gloves, goggles, face

shield, etc.) should be worn when handling infected animals or infectious materials.
(See X. Bloodborne Pathogens; Universal Blood and Body Fluid Precautions)

7. Spills and accidents which result in exposure of people or the environment to infectious materials and/or rDNA molecules are immediately reported to the principal investigator, to the appropriate Department Head, and to the Biosafety Officer. Exposure may require medical evaluation, treatment, and surveillance. Accident investigation may assist in the prevention of similar types of accidents in the future.
8. When it is deemed appropriate by the principal investigator and/or the Biosafety Committee, baseline serum samples for laboratory and other at-risk personnel are collected and stored at the University Health Services. Additional samples may be collected periodically. Serum samples are useful for biological monitoring of workplace exposures in the effort to reduce occupational risks. Stored serum samples are used only to compare pre and post occupational exposure of serum components. Any use of stored samples for any purpose other than those associated with occupational exposures requires the informed consent of the individuals involved.
9. Laboratory personnel are to read and become familiar with the VSU Biosafety Manual and specific standard operating procedures of the laboratory. The principal investigator is responsible for providing supplemental safety training and information for personnel in his/her laboratory.
10. NIH rDNA Guidelines and OSHA Bloodborne Pathogens regulations are examples of two federal regulations requiring appropriate biosafety training for laboratory personnel. Since personnel who are trained and use appropriate biosafety procedures are less likely to lose research time from injuries, providing safety training for personnel is prudent.
11. A high degree of precaution must always be taken with any contaminated sharp items, including needles and syringes, slides, pipettes, capillary tubes, and scalpels. Needles and syringes should be restricted in the laboratory for use only when there is no alternative, such as parenteral injection, phlebotomy, or aspiration of fluids for laboratory animals and diaphragm bottles. Plasticware should be substituted for glassware whenever possible.
 1. Only needle-locking syringes or disposable syringe-needle units (i.e. needle is integral to the syringe) are used for injection or aspiration of infectious materials. Used disposable needles must not be bent, sheared, broken, recapped, removed from disposable syringes, or otherwise manipulated by hand before disposal; rather they must be carefully placed in puncture-resistant containers used for sharps disposal.
 - 2.

BSL-2 Containment Equipment

Biological safety cabinets (BSC) and other appropriate containment devices are to be used whenever laboratory procedures have a good potential for creating aerosols of infectious materials or rDNA molecules. Procedures that may create aerosols include centrifuging, grinding, blending, vigorous shaking or mixing, sonic disruption, opening containers, inoculating animals intranasally, and harvesting tissues from animals or eggs. Biological safety cabinets and other containment devices are to be maintained in good working condition. Certification of biological safety cabinets is to be accomplished annually or whenever the cabinet is moved or the HEPA filter is changed or major repair is accomplished (whenever the contaminated plenum is breached). Certification of biological safety cabinets is conducted by trained personnel and is coordinated by the Biosafety Officer.

BSL-2 Laboratory Facilities

Laboratory facilities are similar to those for BSL-1 with the addition of an autoclave which is readily available and easily accessible. A recording autoclave is required for treating biomedical waste.

Biosafety Level 3 / Laboratory

Biosafety Level 3 is applicable to clinical, diagnostic, teaching, research, or production facilities in which work is done with indigenous or exotic agents which may cause serious or potentially lethal disease as a result of exposure by inhalation. A greater level of attention to microbiological practice, laboratory containment, safety equipment, and facilities is required. All procedures involving the manipulation of infectious materials are conducted within biological safety cabinets or other physical containment devices, by personnel wearing appropriate personal protective clothing and equipment. The laboratory has special engineering and design features. The same standard practices as discussed for BSL-1 are appropriate for BSL-3 with the following additions:

1. Laboratory doors must be kept closed during experiments.
2. Protective clothing is worn with the closed front smock replacing the laboratory coat. All protective clothing is either disposed of in the laboratory or decontaminated by autoclaving prior to laundering.

BSL-3 Special Safety Practices

The same special practices which were appropriate for BSL-2 are appropriate for BSL-3 with the addition of:

1. All activities involving infectious materials or rDNA molecules from BSL-3 organisms are conducted using appropriate containment devices - Biological Safety Cabinets, safety centrifuge cups, etc. No work in open vessels is conducted on the open bench. The significant reduction/prevention of exposure to aerosols is accomplished by a combination of safe work practices and containment equipment.

2.

determine appropriate times and maintain appropriate records of the process.

Autoclaves should receive routine inspection to determine the need for maintenance and repair. Autoclave door gaskets may become distorted if the door is tightly shut for prolonged periods resulting in leaks. Doors should be kept open or loosely closed except when the autoclave serves as a barrier between clean and dirty areas.

Effective decontamination and sterilization by steam depends on the adequacy of circulation of the steam; loads packed tightly may not allow for adequate circulation. The steam must penetrate all packaging materials and contact all surfaces to be decontaminated or sterilized. And, finally the packaging must prevent the recontamination of the sterilized materials. To achieve effective and safe use of the autoclave you must be familiar with and follow your laboratory's procedures regarding:

1. Types of packaging – autoclavable pan, bag in pan, double bag, etc.
2. Separating into pans/bags for autoclaving in the lab
3. Adding water/germicidal solutions - Do not autoclave radioisotopes or explosive or volatile chemicals without checking with radiation safety, laboratory safety and biological safety.
4. Uses of specific autoclaves
5. Proper settings for type of cycle, and type and amount of material. Details of proper operation and settings may be contained in the specific device operation manual. Monitor the autoclave process for proper cycle and length of time. Cycle and time depend on what is being sterilized. For example, liquids would require the use of slow exhaust

Many manipulations of bacterial and viral cultures commonly used in the laboratory generate aerosols of viable organisms. This principle must be remembered when evaluating a person's degree of risk.

-

Biological Safety Cabinets

Primary biohazard containment devices serve to protect laboratory personnel from exposure to infectious aerosols produced by routine procedures. The biological safety cabinet can be an extremely useful containment device for both personnel and product protection. Please see CDC/NIH Primary Containment of Biohazards: Selection, Installation, and Use of Biological Safety Cabinets.

Before purchasing a biological safety cabinet, horizontal flow clean bench or a vertical flow clean bench an approval form must be completed by the investigator and approved by the Biosafety Officer and the Engineering Department of Physical Plant Division

Centrifuges

Centrifuges are an important tool in the microbiological laboratory and must be treated with respect. Each time you use a centrifuge you make a series of choices: Which centrifuge, which rotor, which tubes and adapters, what speed and for how long. In addition, if you are using infectious agents you must decide on the level of containment and then select the appropriate rotor and tubes. Load the infectious agents inside a biological safety cabinet to prevent aerosol exposure. Your choices will affect both your research and yours and others safety.

Always check your user manual for specific requirements as well as load limitations and speed. Operating procedures for each centrifuge must be established by the laboratory supervisor or principle investigator and followed by each operator. These procedures should follow the information provided in the operation manual and guidelines for centrifugation of infectious agents, chemical hazards and/or radioactive materials. Make sure the load is properly balanced – a minor error may not be a problem at low speed but may be serious at higher speeds.

Centrifuge tubes must be selected with the knowledge of the materials they will contain and the pressures they will be under. Plastic centrifuge tubes should be used whenever possible to minimize breakage. Nitrocellulose tubes should only be used when clear, without discoloration, and flexible. It is ad Tw 5.3(ould fol58visisitx01 9pk.00-0.0B2w(a57ocg N2 0 0a06 Tce)]T cen)5.5g.67 0 TD

seal and over the outside of the tube.

Inspect all centrifuge tubes prior to use. Broken, cracked, or damaged tubes are to be discarded. Capped centrifuge tubes should be used whenever possible.

It has been estimated that 80% of centrifuge accidents are operator error. The most common operator errors are failure to secure the rotor to the drive shaft; failure to place lid on the rotor; and failure to secure the lid. Additionally it is very important not to run the rotor above its rated maximum and not to overfill it.

Cryogenic Liquids

Cryogenic liquids are gases that have been transformed into extremely cold refrigerated liquids, which are stored at temperatures below minus 90o C (-130o F). They are normally stored at low pressures in specially constructed multi-walled, vacuum-insulated containers. The hazard potential presented by cryogenic liquids may result from the extreme cold, extreme pressure (which can result from rapid vaporization), and asphyxiation due to the displacement of air.

Appropriate personal protective equipment (heavy leather gloves/gloves for extreme cold, safety shoes, aprons, and eye protection) is to be worn when handling cryogenic liquids or materials preserved in cryogenic liquids.

Lasers

Lasers are tools of biological research and as such must be used with consideration of applicable safety precautions. Refer to the VSU Radiation Safety Officer for appropriate guidance in laser safety.

Ultraviolet Light (UV radiation)

Under certain conditions of radiation intensity and exposure time UV radiation may kill certain types of microorganisms, its greatest effect is against vegetative forms. UV is not a sterilizing agent except in certain exceptional circumstances. It is used to reduce the numbers of microorganisms on surfaces and in the air. The age of the UV lamp, dust accumulations on the bulb, and other factors that impede direct contact of the UV on the microorganisms contribute to decreased efficacy. See Radiation Safety for additional information and safety requirements.

Microwave Ovens

Microwave ovens used in the laboratory may not be used to heat food unless that is the only use of that oven. When melting agar the following precautions must be taken to prevent explosions: Caps on screw-cap bottles must be completely loosened before heating the bottles in the microwave and wear appropriate personal protective equipment including laboratory coat or apron, heat resistant gloves, and face shield.

Laboratory Vacuum Lines

When laboratory vacuum is used to manipulate biohazard materials, suitable filters and traps are to be used to prevent contamination of the vacuum lines and pumps.

Repair and Maintenance of Equipment and Facilities and New Construction

1. University employees or outside vendors undertaking facility expansion, equipment repair and maintenance, and general maintenance activities should not be unnecessarily exposed to biological hazards.
2. It is expected that new construction and renovation projects involving biohazard laboratories are to be reviewed in the planning stages by the University Biosafety Officer, Physical Plant, Campus Planning, and other campus support groups.
3. Repair or routine preventive maintenance of mechanical or laboratory equipment in posted biohazard areas are not to be initiated without prior clearance from the Principal Investigator and Biosafety Officer.
4. Potentially contaminated equipment is not to be removed from biohazard laboratories for repair, servicing, cleaning or to surplus properties or repair shops or other areas until decontamination and removal of biohazard labels have been performed. The investigator or laboratory supervisor is to certify such equipment as being free of biohazard agents. Service personnel may ask laboratory personnel to sign a waiver stating that the piece of equipment has been appropriately decontaminated.
5. Biosafety level 3 agents should not be handled when service personnel are in the laboratory to minimize potential exposure to them.

-

-

X. Vertebrate Animal Biosafety Criteria - Selected Aspects.

-

Specific Medical Concerns for Persons Working with Laboratory Animals.

Allergy and musculoskeletal injury constitute the primary health risks to individuals using and caring for laboratory animals. Allergies are a significant problem, but can be reduced by providing appropriate protective equipment to affected personnel. Musculoskeletal injuries can be minimized by good laboratory planning, use of transport equipment such as carts, and training in lifting techniques and equipment use.

Infectious diseases may constitute a significant risk depending on the species and health status of animals involved, and the level of exposing animal care personnel. Infectious diseases to which animal care personnel may be at risk include a number of viral infections, such as rabies from random source dogs and lymphocytic choriomeningitis from hamsters and mice. In addition to infections potentially acquired from live animals, cell cultures, animal tissues and excreta can

serve as sources of zoonoses. Careful monitoring and quarantine of any animals with potential viral or bacterial infections is a crucial part of quality assurance in animal care programs.

5. All procedures are carefully performed so as to minimize the creation of aerosols.
6. Work surfaces are decontaminated after use or after any spill of viable materials.
7. Doors to animal rooms open inward, are self-closing, and are kept closed when experimental animals are present.
8. All wastes from animal rooms are appropriately decontaminated before disposal. .
9. Infected animal carcasses are incinerated after being transported from the animal room in leakproof, covered containers.
10. An insect and rodent control program is in effect.

Special Practices for Animal BSL 2 may include:

1. Access to animal rooms is limited to personnel who have been advised of potential hazards, meet specific requirements (i.e., immunization, TB tests), and who need to enter the room for program or service work during conduct of the containment phase of operations. In general, persons who are at increased risk of acquiring infection, or for whom infection might be unusually hazardous, are not allowed into the animal room.
2. Under special entry provisions, a hazard warning sign, incorporating the universal biohazard symbol, is posted on the access door to the animal room. The sign should identify the agent(s) in use, names and phone numbers of responsible persons, and special room entry requirements.
3. When appropriate, baseline serum samples from animal care, and other at-risk personnel are collected and stored.
4. Sharp items and accessories, such as needles, syringes, scalpels, glass slides, pipettes, and capillary tubes, are all handled with a high degree of precaution. Such sharp instruments are restricted in the animal facility for use only when there is no alternative, such as for parenteral injection, blood collection, or aspiration of fluids from diaphragm bottles. Plasticware should be used instead of glass whenever possible. The standard operating procedures for sharps should address proper disposal, not recapping of needles, and safe clean up of broken glassware.
5. Cages are properly decontaminated, if appropriate by autoclaving, before they are cleaned and washed.
6. Equipment and work surfaces should be decontaminated on a routine basis, especially after handling the infectious materials and following overt spills or splashes.
7. Incidents involving overt exposures are reported immediately to the lab director and followed by evaluation, surveillance and treatment.

Safety Equipment for Animal Care (primary and secondary barriers)

Laboratory coats, gowns, or uniforms are worn while in the animal room. Gloves are worn when handling infected animals and when skin contact with infectious material is possible. This protective clothing is removed before leaving the animal facility. The animal facility is designed and constructed to facilitate cleaning and housekeeping. Supplies and equipment are stored outside of the animal room area whenever possible. A handwashing sink is available in the room where the infected animals are housed.

Exhaust air is discharged to the outside without being recirculated to other rooms, and it is recommended that the direction of air flow is toward the inside of the room. An autoclave for decontaminating waste should be available in the building with the animal facility.

Biological safety cabinets, other physical containment devices, and/or protective equipment i.e., respirators, face shields, are used whenever procedures with a high potential for creating aerosols are conducted. Such procedures include necropsy of infected animals, harvesting tissues or fluids from infected animals or eggs, intranasal inoculation, and manipulations of high concentrations or large volumes of infectious materials.

*Excerpted or summarized from the CDC/NIH "Biosafety in Microbiological and Biomedical Laboratories" 3rd Edition

-

XI. Floral Biosafety Criteria - Selected Aspects.

-

Biohazard Containment for Plants.

1. The principal purpose of plant containment is to avoid unintentional transmission of recombinant DNA containing plant genome, including nuclear or organelle hereditary material; the release of recombinant DNA derived organisms associated with plants; the release of non-indigenous species; or the release of plant pathogens/pests associated with research at Valdosta State University Facilities.

2. The containment principles used in this section of the biosafety manual are based on the recognition that the organisms to which they apply pose no health threat to human or higher animals unless deliberately modified to do so, and that the intent of containment is to minimize the possibility of unanticipated deleterious effects on organisms and ecosystems outside the experimental facility.

3. The intentional release of genetically engineered organisms and products which are or are believed to be plant pests is regulated under CFR Parts 330 and 340 by the Animal and Plant

Health Inspection Service, United States Department of Agriculture. Biological pesticides and certain field trials are regulated by the United States Environmental Protection Agency. In each case of proposed intentional release, the investigator(s) shall submit appropriate information on anticipated environmental impacts (as submitted to USDA, EPA, or other Federal or State regulatory Agency) for review by the VSU Biosafety Committee.

4. It is the responsibility of the Investigator to obtain any necessary permits for transport and or work with regulated organisms/products. A copy of the permit is to be provided to the office of biosafety.

5. Laboratory experiments with biohazardous plant materials are to be conducted at Biosafety Level 2 (BSL-2) which is a good basic research laboratory and provides the investigator with greater flexibility. BSL-2 is described elsewhere in this manual. Greenhouse containment practices involve a combination of biological and physical protocols.

Greenhouse Biological Containment Practices

Biological containment practices are intended to be used in association with facility design and facility/experimental operational procedures. Effective dissemination of plants by pollen or seed can be prevented by one or more of the following:

1. Preventing insect mediated pollination by appropriate insect control measures within the greenhouse.
2. Covering reproductive structures to prevent pollen dissemination at flowering and seed dissemination at maturity.
3. Removing reproductive structures, employing male sterile strains, or terminating the experiment and harvesting the plant material prior to the reproductive stage.
4. Ensuring that the experimental plants flower at a time of year when non cross-fertile plant is flowering within the normal pollen dispersal range of the experimental plant.
5. Ensuring that no cross-fertile plant is growing within the experimental plant's known pollen dispersal range.

Facilities and definitions.

1. 'Greenhouse' refers to a permanent structure with walls, roof, and floor designed and utilized principally for growing plants in a controlled and protected environment. The walls and roof are usually constructed of transparent or translucent material to allow passage of sunlight for plant growth.
2. 'Greenhouse facility' includes the actual greenhouse rooms or compartments for growing plants plus all immediately contiguous hallways and headhouse areas and is considered part of the confinement area.

XII. Bloodborne Pathogens; Universal Blood and Body Fluid Precautions

The following are the key elements, which can be used at Valdosta State University to control occupational exposures to bloodborne pathogens. All blood and body fluids must be considered as potentially infectious and personnel are to use appropriate protective measures to prevent exposure.

Personnel Practices

Hand-washing:

1. When hands become contaminated with blood or body fluids
2. When gloves are removed
3. Before going to lunch, breaks, or home

Contaminated Needles and Other Sharps:

1. DO NOT recap, bend, or break used needles
2. Discard needles & sharps in appropriate "Sharps" containers
3. Transport reusable sharps in leak-proof puncture-resistant container
4. Use mechanical device (forceps) to place contaminated broken glass into appropriate containers for autoclaving

Personal Protective Equipment for Blood or Body Fluid Contact

1. Gloves when touching blood or body fluids, mucous membranes, or non-intake skin of patients
2. Gloves when handling items or surfaces soiled with blood or body fluids
3. Gloves when performing vascular access procedures (phlebotomy)
4. Appropriate gowns or aprons when splashes or soiling of skin or clothing with blood or body fluids is likely
5. Masks and goggles, or face shield during procedures likely to generate splashes of blood or body fluids into the mouth, nose, or eye

Environmental Controls

General Housekeeping:

1. Maintain work area in clean and sanitary condition
2. Decontaminate work surfaces after procedures and when contaminated
3. Remove any protective work surface coverings when contaminated

Blood or Body Fluid Spills:

1. Soak up spills with absorbent material (paper towels)
2. Decontaminate area with appropriate disinfectant
3. Dispose of contaminated material appropriately

Biomedical Wastes:

1. Are to be disposed of according to State of Georgia Regulations

Transport:

1. Consider all laboratory specimens of human or animal origin as potentially infectious
2. Use leak proof containers for laboratory specimens
3. Place container in a sealable secondary container for transport

Exposures to blood or body fluids via broken skin or needle sticks or mucous membrane contact:

1. Wash affected area immediately and apply first aid
2. Contact VSU Health Service as soon as possible for post exposure follow-up.
3. Report injury to Biosafety Officer.

Biohazard Warning

1. Use appropriate biohazard labels to identify contaminated materials

XIII. Experiments Prohibited at Valdosta State

1. Experiments using pathogenic organisms or DNA from pathogenic organisms classified as requiring Biosafety Level 4 are prohibited.
2. Experiments using any organism or agent that is prohibited by any federal or state agency from importation into Georgia are prohibited.
3. Experiments using agents or organisms that require containment facilities or equipment which are not available at the Valdosta State University are prohibited.

XIV. Transporting Biohazardous Materials

All guidelines of the CDC, NIH, USDA, and EPA are to be followed when biohazardous materials are transported. Individuals transferring **select agents** must adhere to additional regulations specified by the CDC/USDA.

INFORMATION

- CDC Office of Health and Safety Information System <http://www.cdc.gov/od/ohs/>
- USDA, APHIS <http://aphisweb.aphis.usda.gov/>
- Guidelines for the Shipment of Dried Blood Spot Specimens <http://www.cdc.gov/od/ohs/biosfty/driblood.htm>
- Interstate Shipment of Etiologic Agents <http://www.cdc.gov/od/ohs/biosfty/shipregs.htm>

- Packaging and Shipping Instructions <http://www.cdc.gov/od/ohs/biosfty/shipdir.htm>
- USDA, APHIS Forms <http://www.aphis.usda.gov/forms/index.html>

XV. Spills of Biohazards Materials

1. Primary responsibility for preventing or/and containing and cleaning up laboratory spills remains with the principal investigator or laboratory supervisor. Laboratory protocols should be carefully designed to prevent biological, chemical and/or radiation spills.
2. When accidents occur that involve the mishandling or escape of biohazardous materials, the principal investigator or laboratory supervisor is to be notified immediately. Spills of high risk organisms (certain Class 2 and all Class 3) should be reported to the Biosafety Officer during normal working hours or to the Valdosta State University Public Safety Division at the emergency telephone number after normal working hours by the principal investigator or laboratory supervisor. The Public Safety Division will contact the Biosafety Officer for appropriate response. All employees and/or students have an obligation to themselves and their colleagues to report accidents immediately in order to minimize potential hazard.
3. When a biohazardous spill also involves radioactivity, cleanup procedures may have to be modified. The extent of the modification will depend on the level of radiation and the nature of the isotope involved. The Radiation Safety Officer should be called during normal working hours, or the Valdosta State University Public Safety Division should be called after normal working hours.
4. The attached guidelines are intended to assist the principal investigator, laboratory supervisor, and other responsible individuals who may be involved in the cleanup of biological spills.

Biohazardous Spills Inside Laminar Flow Biological Safety Cabinets (LFBSC)

The occurrence of a spill in a biological safety cabinet poses less of a problem than a spill in an open laboratory as long as the spilled materials is contained in the biological safety cabinet. Decontamination of the work zone can usually be effected by direct application of concentrated liquid disinfectants along with a thorough wipe down procedure. Gaseous decontamination may be required to clean up the interior sections of the cabinet.

Procedures for Decontamination of LFBSC

1. Chemical decontamination procedures should be initiated immediately while the biological safety cabinet continues to operate. Continuing the operation of the LFBSC helps to prevent the escape of contaminants from the cabinet.
2. Wearing protective gloves spray or wipe walls, work surfaces, and equipment with an appropriate decontaminating solution. A disinfectant detergent, such as Wescodyne or

Environ has the advantage of detergent action on extraneous organic substances which may interfere with the microbicidal activity of the disinfectant.

3. Flood tray top, drain pans, and catch basins below work surface with decontaminating solution and allow to stand for 20 minutes.
4. Drain excess decontaminating solution from tray and drain pans into cabinet base. Lift out tray and removable exhaust grille work. Clean the top and bottom (underside) surfaces using a sponge or clean cloth soaked in decontaminant solution. Following the cleaning process, replace the tray and grille work in their proper position. Place gloves and sponge or cloth in autoclave pan and autoclave these items.
5. Drain decontaminating solution from cabinet base into appropriate container and

4. Wipe up the spill with the soaked paper towels and place the used towels in an autoclave pan and autoclave.
5. Pour decontaminating solution around and on the area of the spill. Let stand for 20 minutes then wipe up with paper towels. Place gloves and paper towels in autoclave pan

FLOW BIOLOGICAL SAFETY CABINETS (LFBSC)

To be posted near the biosafety cabinet.

1. KEEP THE LFBSC ON.
1. PUT ON PROTECTIVE GLOVES.
1. SPRAY/WIPE WALLS, WORK SURFACES, AND EQUIPMENT WITH DECONTAMINATING SOLUTION.
1. FLOOR TRAY TOP, DRAIN PANS, AND CATCH BASINS WITH DECONTAMINATING SOLUTION.
1. ALLOW TO STAND FOR 20 MINUTES.
1. DRAIN EXCESS SOLUTION INTO CABINET BASE.
1. LIFT OUT TRAY AND REMOVABLE EXHAUST GRILLE WORK.
1. CLEAN TOP AND BOTTOM SURFACES WITH SPONGE/CLOTH SOAKED IN DECONTAMINATING SOLUTION.
1. REPLACE TRAY AND GRILLE WORK.
1. PLACE GLOVES, SPONGE, CLOTH, ETC. IN AUTOCLAVE PAN.
1. DRAIN DECONTAMINATING SOLUTION FROM CABINET BASE INTO AUTOCLAVABLE CONTAINERS.
1. AUTOCLAVE.
1. IF GASEOUS DECONTAMINATION IS NEEDED, CALL THE BIOSAFETY OFFICER.

Checklist

BIOHAZARD SPILL PROCEDURES FOR OUTSIDE LAMINAR

FLOW BIOLOGICAL SAFETY CABINETS (LFBSC)

Minor spills – Class 2 Organisms

To be posted near the biosafety cabinet.

1. WASH HANDS AND OTHER APPARENTLY CONTAMINATED BODY PARTS WITH SOAP AND WATER.
1. POST WARNING TO KEEP NON-ESSENTIAL PERSONNEL FROM SPILL AREA.
1. PUT ON PROTECTIVE GLOVES.
1. COVER SPILL AREA WITH PAPER TOWELS SOAKED IN DECONTAMINATING SOLUTION.
1. WIPE UP SPILL WITH SOAKED PAPER TOWELS.
1. PLACE USED TOWELS IN AUTOCLAVE PAN.
1. POUR DECONTAMINATING SOLUTION AROUND AND ON SPILL AREA.
1. LET SOLUTION STAND FOR 20 MINUTES.
1. WIPE UP WITH PAPER TOWELS.
1. PLACE PAPER TOWELS AND GLOVES IN AUTOCLAVE PAN.
1. WASH HANDS WITH SOAP AND WATER.
1. AUTOCLAVE.

Checklist

BIOHAZARD SPILL PROCEDURES FOR OUTSIDE LAMINAR

FLOW BIOLOGICAL SAFETY CABINET

Major Spills – Class 2 and 3 Organisms

To be posted near the biosafety cabinet.

1. WASH HANDS AND OTHER APPARENTLY CONTAMINATED BODY PARTS

WITH SOAP AND WATER.

1. POST WARNING SIGNS AND CLOSE LABORATORY DOOR.
1. REPORT SPILL TO SUPERVISOR AND BIOSAFETY OFFICER.
1. IF CLOTHING IS CONTAMINATED, REMOVE ALL CONTAMINATED GARMENTS.
1. PLACE CONTAMINATED CLOTHING IN AUTOCLAVE CONTAINER.
1. PUT ON CLEAN GARMENTS.
1. LEAVE LABORATORY FOR 20 MINUTES.
1. CHECK TO SEE THAT LABORATORY DOORS ARE CLOSED AND WARNING SIGNS DISPLAYED UPON RETURNING TO LAB.
1. PUT ON NEEDED SAFETY EQUIPMENT (DISPOSABLE GLOVES, RESPIRATORS, ETC.).
1. PLACE PAPER TOWELS SOAKED IN DECONTAMINATION SOLUTION OVER THE SPILL.
1. POUR DECONTAMINATION SOLUTION AROUND SPILL – ALLOW SOLUTION TO FLOW INTO SPILL. DO NOT POUR DECONTAMINATION SOLUTION INTO SPILL.
1. LET STAND FOR AT LEAST 20 MINUTES.
1. TRANSFER CONTAMINATED MATERIALS TO AUTOCLAVE CONTAINER USING AUTOCLAVABLE DUST PAN AND SQUEEGEE.
1. PLACE DUST PAN AND SQUEEGEE IN AUTOCLAVE CONTAINER.
1. REMOVE GLOVES AND OTHER CONTAMINATED GARMENTS AND PLACE IN AUTOCLAVE CONTAINER.
1. WASH FACE, HANDS, AND OTHER APPARENTLY CONTAMINATED BODY PARTS.
1. AUTOCLAVE ALL MATERIALS THAT REQUIRE AUTOCLAVING.

Liquid Disinfectants

Laboratory personnel should be familiar with the various disinfectants that will effectively kill the biohazardous agents being used. The following information is provided to assist in your selection of appropriate disinfectants.

Alcohols – Ethyl and Isopropyl are good disinfectants for the vegetative forms of bacteria and enveloped viruses.

Ethyl Alcohol

1. Use Dilution: 70-95%
2. Inactivates: vegetative bacteria and enveloped viruses, has variable results with non-enveloped
3. viruses and is ineffective with bacterial spores.
4. Other Characteristics: flammable, eye irritant, and toxic [Threshold limit value (TLV) – 1000 ppm]

Isopropyl Alcohol

the compound in which it is contained. Allow a contact time of 10 to 30 minutes.

1. Use Dilution: 25 to 1600 ppm of available iodine. Solutions containing 75 to 150 ppm are generally recommended.
2. Inactivates: vegetative bacteria, fungi and viruses. There is poor activity against bacterial spores.
3. Other Characteristics: Although iodophors are less harmful to man than chlorine compounds they can irritate the skin and eyes. Iodophors are corrosive (less than chlorine), they leave a residue and may stain. Iodophor stains, however, can be readily removed with solutions of sodium thiosulfate ($\text{Na}_2\text{S}_2\text{O}_3$). As with the chlorine compounds, iodophors are rapidly inactivated by organic matter. One advantage is that iodophors have a built-in indicator. As long as the solution is brown or yellow it is still active.

Phenolic Compounds – These are effective against vegetative bacteria (including *Mycobacterium*

tuberculosis), fungi, and enveloped viruses. Effectiveness against non-enveloped viruses is variable depending on the virus. The phenols are ineffective against bacterial spores.

1. Use Dilutions: 1.0 – 5.0% Solutions containing 0.5 – 2.0% phenol are effective against enveloped viruses.
2. Other Characteristics: Phenols are corrosive and may leave a sticky, gummy residue.
3. Phenolic compounds are irritating to the skin and eyes and are relatively toxic – Phenol
4. TLV for skin is 5 ppm.

Quaternary Ammonium Compounds – The efficacy of Quaternary Ammonium compounds still generates considerable controversy. Quats are effective in destroying ordinary vegetative bacteria and lipid containing virus but are not effective against *Pseudomonas*, *Proteus* and other gram-negative bacilli. Also, Quats are not effective against bacterial spores at the usual use concentrations of 1:750.

1. Use Dilutions: 0.1 to 2.0%
2. Other Characteristics: Quats are surface-active compounds which possess the useful property of lowering the surface tension of the solution.
3. Other advantages include being nontoxic, odorless, nonstaining, noncorrosive to metals and stable. If used at recommended concentrations, Quats are nonirritating.
4. Quaternary Ammonium compounds are rapidly inactivated by organic matter.

BUILDING & ROOM NO(s) _____

GRANTING AGENCY _____

GRANT NO. (IF APPLICABLE) _____

TITLE OF GRANT OR PROJECT:

A. Describe the experiment involving recombinant DNA techniques. Your description is to be sufficiently complete so as to provide committee members an understanding of what you intend to do and how you will do it. A summary or abstract of your methods and materials section will also be helpful. Please reference this discussion to appropriate NIH Guidelines and/or USDA/APHIS, and EPA regulations.

-Page 2-

MEMORANDUM OF UNDERSTANDING AND AGREEMENT

FOR RECOMBINANT DNA EXPERIMENTS (Continued)

B. ASSESSMENT LEVELS OF PHYSICAL AND BIOLOGICAL CONTAINMENT.

1. Describe how you intend to meet physical and biological containment requirements (reference NIH/USDA/EPA guidelines).
2. Will this project involve environmental release?
3. Describe procedures and precautions to be followed in transporting biohazardous agents between laboratories.

C. Agreements

___ I agree to accept responsibility for training of all laboratory workers involved in the project.

___ I agree to comply with all appropriate requirements pertaining to shipment and transfer of recombinant DNA materials.

VALDOSTA STATE UNIVERSITY

MEMORANDUM OF UNDERSTANDING AND

AGREEMENT (MUA) FOR BIOHAZARDS OTHER THAN RECOMBINANT DNA EXPERIMENTS

DATE: _____

RESEARCHER'S
NAME _____

RESEARCHER'S TITLE _____

PHONE NO. _____

DEPARTMENT _____

BUILDING. & ROOM NO(s) _____

GRANTING AGENCY _____

GRANT NO. (IF APPLICABLE) _____

TITLE OF GRANT OR PROJECT:

A. Describe the experiments involving biohazard(s). Your description is to be sufficiently complete so as to provide committee members an understanding of what you intend to do and how you will do it.

(CONTINUED)

B. Assess the levels of physical containment required for the experiments.

C. Describe the facilities and specific procedures that will be used to provide the required levels of containment.

D. Describe the procedures and precautions to be followed if biohazardous organisms or agents are to be transported between laboratories.

-Page 2-

**MUA FOR BIOHAZARDS OTHER THAN RECOMBINANT DNA EXPERIMENTS
(CONTIUED)**

E. The undersigned agree to certify the following conditions of the proposed research:

1.

individuals listed by name, before commencing the project described in this MUA:

G. We certify that the Valdosta State University Committee on Biosafety has reviewed the proposed project and has found it to be in compliance with the VSU Biosafety Manual, which outlines standards for conducting experiments with biohazardous agents.

Chairperson, VSU Committee on Biosafety

DATE

VSU Biosafety Officer

DATE

VALDOSTA STATE UNIVERSITY

BIOSAFETY INFECTIOUS AGENT RISK ASSESSMENT

Investigator Name _____

Department _____

Section I - Agent Identification:

Agent Name: (common) _____

(Scientific) _____

Nature of Agent ___bacterial ___viral ___parasitic ___fungal
___other

Supplier/Source:

OUTSIDE

Name

—

Address

Phone (_____) - _____

Other
information _____
-

DESCRIBE PRIMARY ISOLATION

Section II - Biosafety level

Biosafety Level _____1 _____2 _____3 _____4

Source of determination

-Page 2-

VALDOSTA STATE UNIVERSITY

BIOSAFETY INFECTIOUS AGENT RISK ASSESSMENT (CONTINUED)

Describe Unique features of the agent (if any) -

Section III - Health Hazard Information

_____ Primary pathogen _____ Opportunistic pathogen

Name of Disease/Illness _____

Describe Signs and Symptoms _____

Incubation Period _____

Method of Transmission

Direct Transmission _____yes _____no _____unknown

Indirect Transmission _____yes _____no _____unknown

Airborne Transmission _____yes _____no _____unknown

Direct contact _____yes _____no _____unknown

Vector borne _____yes _____no _____unknown

Droplet _____yes _____no _____unknown

Vehicle borne _____yes _____no _____unknown

Aerosol _____yes _____no _____unknown

Others_____

Lab coat/gown/apron _____yes _____no

Respirator (HEPA filter) needed _____yes _____no

If yes,
describe_____

-Page 4-

VALDOSTA STATE UNIVERSITY

BIOSAFETY INFECTIOUS AGENT RISK ASSESSMENT (CONTINUED)

Section V - Background Information:

Oxygen requirements _____

Gram stain_____

Spore forming _____Yes _____No

Generation
time_____

Incubation
period_____

Nutrient requirements

Sensitivities:

Desiccation _____yes _____no

Light _____yes _____no

Temperature _____yes _____no

Chemicals _____yes _____no

Specify _____

Radiation _____yes _____no

Antibiotic _____yes _____no

Specify _____

Environmental Factors: (extrinsic factors that affect agent and opportunity for spread)

Biosafety Committee Assessment _____Agree _____Disagree

Modifications _____

VALDOSTA STATE UNIVERSITY

BIOSAFETY INFECTIOUS AGENT RISK ASSESSMENT (CONTINUED)

Date _____

Principal

Investigator _____

Department _____

Telephone _____

Building & Room Number where unit is to be installed _____

Equipment List/Desired Unit

_____ Class II A Biological Safety Cabinet

_____ Class II B1 Biological Safety Cabinet

_____ Class II B2 Biological Safety Cabinet

_____ Glove Box

_____ Horizontal Flow Clean Bench

_____ Vertical Flow Clean Bench

The above unit is for use with:

_____ Tissue Culture
(Identify) _____

_____ Clinical Specimens
(Identify) _____

_____ Pathogenic Microorganisms
(Identify) _____

CDC/Biosafety Level _____2 _____3 _____4

_____ Carcinogens
(Identify) _____

Amounts to be used _____

_____ Media Preparation

_____ Sterile Apparatus Assembly

_____ Flammable or/and Toxic Chemicals

_____ Other

Will unit be attached to building exhaust system? _____yes _____no

This Unit Approved By:

Chairperson, Biosafety Committee

Date

Engineering Dept./Physical Plant

Date

One of these forms must be filed with Procurement for each individual unit purchased.

VSU BIOSAFETY

LABORATORY EQUIPMENT APPROVAL FORM

Date _____

Principal
Investigator _____

Department _____
Telephone _____

Building & Room Number where unit is to be
installed _____

Equipment List/Desired Unit

_____ Class II A Biological Safety Cabinet

_____ Class II B1 Biological Safety Cabinet

_____ Class II B2 Biological Safety Cabinet

_____ Glove Box

_____ Horizontal Flow Clean Bench

_____ Vertical Flow Clean Bench

The above unit is for use with:

_____ Tissue Culture

(Identify) _____

_____ Clinical Specimens

(Identify) _____

_____ Pathogenic Microorganisms

(Identify) _____

CDC/Biosafety Level _____ 2 _____ 3 _____ 4

_____ Carcinogens

(Identify) _____

Amounts to be
used _____

_____ Media Preparation

_____ Sterile Apparatus Assembly

_____ Flammable or/and Toxic Chemicals

_____ Other

Chairperson, Biosafety Committee

Date

Engineering Dept./Physical Plant

Date

One of these forms must be filed with Procurement for each individual unit purchased.

VALDOSTA STATE UNIVERSITY - BIOHAZARDOUS RESEARCH CHECKLIST

DATE _____

INVESTIGATOR _____ PHONE

NO. _____

DEPARTMENT _____

LOCATION _____

CO-PIs _____

FUNDING AGENCY _____

GRANT: NEW _____
CONTINUATION/RENEWAL _____

MUA PREVIOUSLY SUBMITTED--GIVE APPROVAL DATE _____

TITLE OF PROPOSAL _____

PLEASE INDICATE WHETHER THE STUDIES IN THE ACCOMPANYING RESEARCH GRANT PROPOSAL INVOLVE ANY OF THE FOLLOWING BY COMPLETING THE APPROPRIATE SPACES BELOW. (FOR FURTHER INFORMATION, SEE THE VSU BIOSAFETY MANUAL.)

TYPE I. In vitro construction and/or propagation of recombinant DNA molecules

_____yes _____no

A. Formation of rDNAs containing genes for the biosynthesis of toxic molecules lethal for vertebrates at an LK50 of less than 100 ng per kg body weight

_____yes _____no

B. Deliberate release into the environment of any organism containing rDNA

_____yes _____no

C. Deliberate transfer of a drug-resistance trait in microorganisms not known to acquire it naturally.

_____yes _____no

D. Involves plant pests

_____yes _____no

E. Involves transformation of whole plants

_____yes _____no

F. Involves transformation of animals

_____yes _____no

TYPE II. Experiments with organisms of demonstrated human pathogenicity. Specify agent(s) in subsequent section.

A. Involves the infection of animals

_____yes _____no

TYPE III. Experiments on oncogenic (cancer-causing) animal viruses. Specify viruses in subsequent section.

A. Involves the infection of animals

_____yes _____no

TYPE 25 -2.3 TD0 TD()8.7olno

TYPE VI. Experiments involving the movement into Georgia of:

A. Pathogens that adversely affect plants or animals

_____yes _____no

B. Any non-indigenous species of plants or animals (including offspring).

_____yes _____no

C. Any indigenous species of plants or animals infected with pathogenic organisms outside of the state

_____yes _____no

TYPE VII. Experiments involving the transfer or receipt of select agents as defined in the Code of Federal Regulations (42 CFR Part 72.6). Specify agents in subsequent section.

_____yes _____no

Type II. Specify agents:

Type III. Specify viruses:

Type IV Specify cell lines/types of tissue cultures:

Type V Specify organisms:

Type VI Specify select agents:

Signature of Investigator

Date

Signature of Department Head

Date

Submit this form to the VSU Biosafety Officer